

6. Ekstrom P, et al. Prevention of peptic ulcer and dyspeptic symptoms with Omeprazole in patients receiving Nonsteroidal Anti-inflammatory Drug continuous therapy. A nordic multicentre study. *Scand J Gastroenterol* 1996; 31: 753-8.
7. Feldman M. Peptic ulcer disease. In: Dale DC, Federman DD eds. *Scientific American Medicine, Section 4, Gastroenterology II*, 2000: 2-3.
8. Graham DY, et al. Ulcer prevention in long-term users of Nonsteroidal Antinflammatory Drugs. *Arch Intern Med* 2002; 162: 169 – 175.
9. Graham DY. Critical effect of *Helicobacter pylori* infection on the effectiveness of omeprazole for prevention of gastric or duodenal ulcers among chronic NSAID users. *Helicobacter* 2002; 7: 1-8.
10. Graham DY. NSAIDS, *Helicobacter pylori* and Pandora Box. *N Engl. J. Med.* 2002; 347: 2162-2164.
11. Hawkey CJ, et al. Omeprazole compared with Misoprostol for ulcers associated either Nonsteroidal Anti-inflammatory Drugs. *N Engl J Med* 1998; 338: 727-34.
12. Jia-Qing, et al. Role of *Helicobacter pylori* infection and non-steroidal anti-inflammatory drugs in peptic ulcer disease : a meta-analysis. *Lancet* 2002; 359:14-22.
13. Kelly YP, et al. Risk of aspirin – associated major upper gastrointestinal bleeding with enteric-coated or buffered products. *Lancet* 1996; 384: 1413-6.
14. Koch M, et al. Prevention of Nonsteroidal Anti-inflammatory drugs-induced gastrointestinal mucosal injury. A meta-analysis of randomised controlled clinical trial. *Arch Intern Med* 1996; 156: 2321-32.
15. Lanza FL. A guideline for the treatment and prevention of NSAID-induced ulcers. *Am J Gastroenterol* 1998; 93: 2037-46.
16. Silverstein, et al. Misoprostol reduces serious gastrointestinal complications in patients with rheumatoid arthritis receiving Nonsteroidal Anti-inflammatory drugs. A. randomised, double-blind, placebo-controlled trial. *Ann Intern Med* 1995; 123: 241-9.
17. Wolfe MM, et al. Gastrointestinal toxicity of Non-steroidal Anti-inflammatory Drugs. *N Engl J Med* 1999; 340: 1888-99.
18. Yeomans ND, et al. A comparison of Omeprazole with Ranitidine for ulcers associated with Nonsteroidal Anti-inflammatory Drugs. *N Engl J Med* 1998; 338:719-26.
19. Chan FKL et al. Combination of a cyclo-oxygenase-2 inhibitor and a proton-pump inhibitor for prevention of recurrent ulcer bleeding in patients at very high risk: a double-blind, randomised trial. *Lancet.* 2007; 369: 1621-6.
20. Heidelbaugh JJ, Goldberg KL, Inadomi JM. Overutilization of proton pump inhibitors: a review of cost-effectiveness and risk [corrected] *Am J Gastroenterol.* 2009 Mar; 104:27-32.
21. Lanza FL, Chan FK, Quigley EM. Practice Parameters Committee of the American College of Gastroenterology. Guidelines for prevention of NSAID-related ulcer complications. *Am J Gastroenterol.* 2009;104:728-38.
22. Schlansky B, Hwang JH. Prevention of nonsteroidal anti-inflammatory drug-induced gastropathy. *J Gastroenterol.* 2009; 19:44-52.
23. Gilard M, Arnaud B, Cornily JC, Le Gal G, Lacut K, Le Calvez G, et al. Influence of omeprazole on the antiplatelet action of clopidogrel associated with aspirin. The randomized, double-blind OCLA (Omeprazole CLopidogrel Aspirin) study. *J Am Coll Cardiol* 2008;51(3):256-60.

